Noname manuscript No.
(will be inserted by the editor)

Low Temperature Transport Properties of Very Dilute Classical Solutions of ^{3}He in Superfluid ^{4}He

Gordon Baym · D. H. Beck · C. J. Pethick

arXiv:1408.1619v1 [cond-mat.quant-gas] 7 Aug 2014

Received: date / Accepted: date

Abstract We report microscopic calculations of the thermal conductivity, diffusion constant and thermal diffusion constant for classical solutions of ^{3}He in superfluid ^{4}He at temperatures $T \lesssim 0.6$ K, where phonons are the dominant excitations of the ^{4}He. We focus on solutions with ^{3}He concentrations $\lesssim 10^{-3}$, for which the main scattering mechanisms are phonon-phonon scattering via 3-phonon Landau and Beliaev processes, which maintain the phonons in a drifting equilibrium distribution, and the slower process of ^{3}He-phonon scattering, which is crucial for determining the ^{3}He distribution function in transport. We use the fact that the relative changes in the energy and momentum of a ^{3}He atom in a collision with a phonon are small to derive a Fokker-Planck equation for the ^{3}He distribution function, which we show has an analytical solution in terms of Sonine polynomials. We also calculate the corrections to the Fokker-Planck results for the transport coefficients.

This research was supported in part by NSF Grants PHY08-55569, PHY09-69790, and PHY13-05891. Author GB is grateful to the Aspen Center for Physics, supported in part by NSF Grant PHY-1066292, and the Niels Bohr International Academy, where parts of this research were carried out.

Gordon Baym
Department of Physics, University of Illinois, 1110 W. Green Street, Urbana, IL 61801, USA, and
The Niels Bohr International Academy, The Niels Bohr Institute, University of Copenhagen, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark
E-mail: gbaym@illinois.edu

D. H. Beck
Department of Physics, University of Illinois, 1110 W. Green Street, Urbana, IL 61801, USA

C. J. Pethick
Department of Physics, University of Illinois, 1110 W. Green Street, Urbana, IL 61801, USA,
The Niels Bohr International Academy, The Niels Bohr Institute, University of Copenhagen, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark, and
NORDITA, Roslagstullsbacken 23, SE-10691 Stockholm, Sweden

Keywords dilute solutions of ^{3}He in ^{4}He · transport · diffusion · thermal conductivity · Boltzmann equation · Fokker-Planck equation · neutron electric dipole moment experiment

PACS 67.60.G- , 13.40.Em · 05.20.Dd

August 8, 2014

1 Introduction

Problems worthy of attack, prove their worth by hitting back [1]

Historically, the study of equilibrium and transport processes in the helium liquids revealed important information about the basic physics of quantum liquids [2]. A proposed experiment at Oak Ridge National Laboratory [3] to search sensitively for the electric dipole moment of the neutron has renewed interest in the transport properties of dilute solutions of ^{3}He in liquid ^{4}He in low concentration regimes never before studied in detail. The experiment will look for the effect of an electric field on the capture rate of polarized neutrons on polarized ^{3}He atoms dissolved in ^{4}He.

In a relative spin singlet state, the capture rate of a neutron on a ^{3}He atom can proceed through a virtual α particle state, and is thus enormous compared with the rate for a relative triplet state. The experiment therefore aims to detect the precession of the neutron in an electric field from a change in the capture rate. With time, however, the ^{3}He atoms depolarize through scattering against the container walls, and in the course of the experiment they will be driven out of the experimental volume by a phonon wind generated by a heater in the liquid ^{4}He, and replaced with fresh polarized ^{3}He.

A novel pilot experiment was performed by Lamoreaux et al. [4], who measured the ^{3}He density gradient induced by a heat flow at temperatures $\lesssim 0.6$ K, at which the effects of rotons are negligible. In this experiment, the ^{3}He number concentration, $x_3 = n_3/(n_3 + n_4)$, where n_3 and n_4 are the ^{3}He and ^{4}He number densities, lay in the range 7×10^{-5} to 1.5×10^{-3} in the non-degenerate regime. The experiment was initially interpreted in terms of diffusion of ^{3}He atoms with respect to a gas of phonons [4,5], but in Ref. [6] we showed that the relevant transport coefficient is the total thermal conductivity of the liquid, which consists of two contributions, one from the ^{3}He and another from diffusive flow of phonons relative to ^{3}He. This latter diffusion does not necessarily involve net relative motion of ^{4}He atoms relative to ^{3}He, as does diffusion in a normal system, since the net flux of ^{4}He atoms has a contribution from the superfluid as well as that from the phonons.

In transport in this relatively high-x_3 regime, the ^{3}He-^{3}He interactions are sufficiently strong that they keep the ^{3}He in thermal equilibrium at rest at the local temperature $T(\mathbf{r})$, while phonon-phonon interactions keep the phonons in drifting local equilibrium. In thermal transport, phonons transfer momentum to the ^{3}He atoms, via the ^{3}He-phonon interactions. The calculations in

Ref. [6] took into account a number of physical effects not included in earlier calculations of the thermal conductivity. Among these are phonon-phonon scattering, which rapidly establishes thermal equilibrium among phonons traveling in a given direction, ^{3}He-^{3}He scattering, which is effective in maintaining thermal equilibrium of these atoms, and energy transfer in ^{3}He-phonon collisions, which reduces the rate of these collisions. These calculations agree well with the data of Ref. [4].

By contrast, in the proposed search for the neutron electric dipole moment, the concentration of ^{3}He will be in the much lower range 10^{-9}-10^{-11}. Here thermal conduction is essentially limited by transfer of phonon momentum to the container walls by viscous forces, with the ^{3}He playing a negligible role. Furthermore, collisions of ^{3}He with the phonons drive the ^{3}He away from equilibrium.

In this paper, starting from microscopic scattering processes we calculate transport coefficients of dilute solutions in the regime where phonons dominate the ^{4}He excitations and the ^{3}He concentrations are low enough for classical statistics to be valid. This regime encompasses the range at relatively high ^{3}He concentrations where measurements have been carried out [4,7], to the range of natural concentration, $x_3 \sim 10^{-6}$ where future measurements are being prepared [8], to the low concentrations anticipated in the Oak Ridge neutron electric dipole moment experiment [3]. Our primary focus here is on the processes of thermal conduction and diffusion.

In the high-x_3 regime, ^{3}He-^{3}He collisions maintain the ^{3}He quasiparticles in a drifting equilibrium distribution. For this case as we showed in Ref. [6], in calculating the leading low-temperature contributions to the transport coefficients, one may neglect the energy transfer in ^{3}He-phonon collisions. At low x_3, however, the effects of ^{3}He-^{3}He collisions are negligible and as we show, it is necessary to take into account the energy transfer even in calculating the leading low-temperature properties. The reason for this is that while the fractional changes in the momentum and energy of a ^{3}He quasiparticle in scattering by a phonon are both small, they are of the same order, $\sim k/p \sim (T/m^*s^2)^{1/2}$, where $\mathbf{k}$ is the momentum transfer, p the momentum of a quasiparticle, m^* the ^{3}He effective mass, and s the sound velocity in ^{4}He. To calculate the leading low-temperature behavior we expand the collision term in the ^{3}He Boltzmann equation to leading non-trivial (second) order in k/p and derive a Fokker-Planck equation for the distribution function. Quite remarkably, we find that the eigenfunctions of the collision integral may be found analytically and have the form of Sonine polynomials, which are familiar in the theory of transport coefficients in classical, single-component gases [9,11].

We begin in Sec. 2 by reviewing the conservation laws obeyed by the phonons and ^{3}He, and define the particle and heat currents in detail. This section follows in large part the general approach to the dissipative hydrodynamics of mixtures in Ref. [10], but with crucial differences. Then in Sec. 3, we review the microscopic ^{3}He-^{3}He interactions, ^{3}He-phonon interactions, and phonon-phonon interactions underlying the transport. Here we do not consider effects, such as phonon scattering from the walls, whose role depends on

the specific geometry in question. The effects of walls, which are critical at ultralow x_3 as will be encountered in the nEDM experiment, will be discussed in Ref. [12]. The scattering of phonons on ^{3}He is characteristically that of a massless particle, as in Rayleigh scattering. We then develop the ^{3}He and phonon Boltzmann equations in Secs. 4 and 5, and discuss recoil corrections in Sec. 6. The resulting transport coefficients are presented in Sec. 7. In general we work in the linear approximation, assuming that the deviations from equilibrium are small.

2 Hydrodynamics and Conservation Laws

In dilute solutions of ^{3}He in low temperature ^{4}He in bulk, mean free paths are generally sufficiently small that the system can be described in terms of dissipative hydrodynamics. In addition, at low temperatures the dominant bosonic excitations of solutions are phonons. We begin by laying out the general framework of the hydrodynamics, explicitly in terms of the ^{3}He and ^{4}He number densities, the superfluid velocity, and the normal fluid ^{3}He and phonon velocities. In a later section we calculate the needed transport coefficients. The description we give here parallels in many ways the description of a normal two-component fluid in Ref. [10], as extended to dilute solutions by Khalatnikov and Zharkov [13,14], with the identification of the phonons and the ^{3}He as the two fluids. An important difference from two component normal fluids is that here the excitations of the ^{4}He are not conserved, and furthermore a phonon drift with respect to the ^{3}He is a heat flow, whereas in a two-component normal fluid, heat flow is energy transport in the absence of relative motion of the two components.

2.1 Conservation of particle number and momentum

The ^{3}He excitation spectrum is effectively

$$\epsilon_p = \epsilon_0 + p^2/2m^*, \tag{1}$$

where the ^{3}He effective mass is $m^* = m_3 + \delta m \simeq 2.34 m_3$ [15]; the δm arises from backflow in the ^{4}He as a ^{3}He moves through it. The mean field shift ϵ_0 depends on both the ^{3}He and ^{4}He densities; however at low concentrations the former dependence is negligible, and $\partial\epsilon_0/\partial n_4 = (m_4 s^2/n_4)(1+\alpha)$ where $\alpha \simeq 0.28$ is the fractional excess molar volume of the ^{3}He. At the temperatures and concentrations of interest, the ^{3}He atoms obey Boltzmann statistics, with the equilibrium density given by

$$n_3 = \nu \int \frac{d^3p}{(2\pi)^3} e^{-\beta(p^2/2m^* + \epsilon_0 - \mu_3)} = \nu e^{\beta(\mu_3 - \epsilon_0)}/\lambda_{th}^3, \tag{2}$$

where μ_3 is the ^{3}He chemical potential, ν is the number of spin degrees of freedom: $\nu = 1$ in a fully spin-polarized gas and 2 in an unpolarized gas,

$\beta = 1/T$, and $\lambda_{th} = \sqrt{2\pi/m^*T}$ is the ^{3}He thermal wavelength. We generally work in units with $\hbar = k_B = 1$. At the low concentrations of interest, effects of ^{3}He-^{3}He interactions on the thermodynamics are negligible.

The conservation law for ^{3}He atoms is

$$\frac{\partial n_3}{\partial t} + \boldsymbol{\nabla} \cdot (n_3 \mathbf{v}_3) = 0, \tag{3}$$

while the conservation law for ^{4}He is

$$m_4 \frac{\partial n_4}{\partial t} + \boldsymbol{\nabla} \cdot \mathbf{g}_4 = 0, \tag{4}$$

where $\mathbf{v}_3$ is the ^{3}He flow velocity, and $\mathbf{g}_4 = \rho_s \mathbf{v}_s + \delta m\, n_3 \mathbf{v}_3 + \rho_{ph} \mathbf{v}_{ph}$ is the ^{4}He mass current, with $\mathbf{v}_s$ the superfluid velocity, $\mathbf{v}_{ph}$ the phonon (normal) fluid velocity, and m_4 the ^{4}He atomic mass. The ^{4}He mass density is $\rho_4 = m_4 n_4$. The superfluid mass density is $\rho_s = \rho_4 - \rho_{ph} - \delta m\, n_3$, since the ^{3}He effective mass correction does not participate in superfluid flow. The phonon fluid velocity is defined by writing the momentum density carried by phonons as $\rho_{ph} \mathbf{v}_{ph}$, with the normal mass density of the phonons given by

$$\rho_{ph} = \beta \int \frac{d^3q}{(2\pi)^3} \frac{q^2}{3} n_q^0 (1 + n_q^0) = \frac{2\pi^2}{45} \frac{T^4}{s^5}, \tag{5}$$

where s is the first sound velocity, $\mathbf{q}$ the phonon momentum, and $n_q^0 = \left(e^{\beta s q} - 1\right)^{-1}$.

The mass current, or total momentum density, $\mathbf{g}$, in the solution is given by

$$\mathbf{g} = \rho_s \mathbf{v}_s + \rho_{ph} \mathbf{v}_{ph} + m^*\, n_3 \mathbf{v}_3 = \mathbf{g}_4 + \mathbf{g}_3^0, \tag{6}$$

where $\mathbf{g}_3^0 = m_3 n_3 \mathbf{v}_3$ is the momentum density carried by ^{3}He atoms, not the fully dressed quasiparticles. In the absence of ^{4}He and ^{3}He mass flow, $\mathbf{v}_s = -(\rho_{ph}/\rho_s)\mathbf{v}_{ph}$; at the temperatures and concentrations of interest, $v_s \ll v_{ph}$.

The linearized superfluid acceleration equation is

$$m_4 \frac{\partial \mathbf{v}_s}{\partial t} + \boldsymbol{\nabla} \mu_4 = 0, \tag{7}$$

where μ_4 is the ^{4}He chemical potential; note that in the absence of superfluid acceleration, μ_4 is independent of position. In addition the hydrodynamic equation for conservation of total momentum in the dilute solutions, is, in linear order,

$$\frac{\partial \mathbf{g}}{\partial t} + \boldsymbol{\nabla} P = \mathbf{H}, \tag{8}$$

where P is the total pressure and $\mathbf{H}$ is the gradient of the viscous stress tensor. In the phonon regime, the phonon first viscosity is dominant; indeed the phonon contributions to the dissipative second viscosity terms in Eqs. (7) and (8) vanish [17]. Similarly, the viscosity of the ^{3}He gas can be neglected,

as can small second viscosity effects associated with heat transfer between the ^{3}He and phonons. Then

$$\mathbf{H} \equiv \eta_{ph} \left(\nabla^2 \mathbf{v}_{ph} + \frac{1}{3} \boldsymbol{\nabla} (\boldsymbol{\nabla} \cdot \mathbf{v}_{ph}) \right), \tag{9}$$

where η_{ph} is the first viscosity of the phonon fluid.

The ^{3}He momentum density obeys [18]

$$\frac{\partial}{\partial t} \left(\mathbf{g}_3 - \delta m n_3 \mathbf{v}_s \right) + n_3 \boldsymbol{\nabla} \mu_3 + S_3 \boldsymbol{\nabla} T = \frac{\partial \mathbf{g}_3}{\partial t} \Big|_{coll}, \tag{10}$$

where $\mathbf{g}_3$ is the momentum density of the dressed ^{3}He quasiparticles,

$$S_3 = \left(\frac{5}{2} - \beta(\mu_3 - \epsilon_0) \right) n_3 = \left[\frac{5}{2} - \ln \left(\frac{n_3 \lambda_{th}^3}{\nu} \right) \right] n_3. \tag{11}$$

is the ^{3}He entropy density, and $(\partial \mathbf{g}_3/\partial t)|_{coll}$ is the rate of change of the ^{3}He momentum density resulting from collisions with phonons. This equation can also be written in terms of the ^{3}He partial pressure, $P_3 = n_3 T$, if one uses the relation $dP_3 = S_3 dT + n_3 d(\mu_3 - \epsilon_0)$.

Similarly the phonon momentum density obeys

$$\frac{\partial}{\partial t} (\rho_{ph} (\mathbf{v}_{ph} - \mathbf{v}_s)) + S_{ph} \boldsymbol{\nabla} T = - \frac{\partial \mathbf{g}_3}{\partial t} \Big|_{coll} + \mathbf{H}, \tag{12}$$

where

$$S_{ph} = s^2 \rho_{ph} / T == \frac{2\pi^2}{45} \left(\frac{T}{s} \right)^3 \tag{13}$$

is the ^{4}He entropy density; this equation can similarly be written in terms of the phonon partial pressure, $P_{ph} = T S_{ph}/4$ if one uses the relation $dP_{ph} = S_{ph} dT - \frac{3}{4} s \rho_{ph} ds$.

With the Gibbs-Duhem relation,

$$\boldsymbol{\nabla} P = n_4 \boldsymbol{\nabla} \mu_4 + n_3 \boldsymbol{\nabla} \mu_3 + S \boldsymbol{\nabla} T, \tag{14}$$

conservation of momentum implies

$$\rho_{ph} \frac{\partial \mathbf{v}_{ph}}{\partial t} + m^* n_3 \frac{\partial \mathbf{v}_3}{\partial t} + n_3 \boldsymbol{\nabla} \mu_3 + \frac{\rho_{ph} + \delta m \, n_3}{m_4} \boldsymbol{\nabla} \mu_4 + S \boldsymbol{\nabla} T = \mathbf{H}; \tag{15}$$

here S is the total entropy density. equal at low temperatures to $S_{ph} + S_3$. In a steady state,

$$n_3 \boldsymbol{\nabla} \mu_3 + S \boldsymbol{\nabla} T = \mathbf{H}; \tag{16}$$

when the phonon viscosity contribution is negligible, a condition realized at ^{3}He relative concentrations $\gtrsim 10^{-4}$, a temperature gradient inevitably accompanied by a ^{3}He chemical potential gradient and vice versa.

2.2 Energy conservation

Using Eqs. (3)-(15), together with the relation for the internal energy $dE_{int} = TdS + \mu_4 dn_4 + \mu_3 dn_3$, we readily find the equation for conservation of energy,

$$\frac{\partial E}{\partial t} + \boldsymbol{\nabla} \cdot \mathbf{j}_E = 0, . \tag{17}$$

where the energy current is

$$\mathbf{j}_E = \mathbf{g}_4 \mu_4 / m_4 + n_3 \mathbf{v}_3 \mu_3 + T \mathbf{v}_3 S_3 + T \mathbf{v}_{ph} S_{ph} + \mathbf{Q}_3. \tag{18}$$

(For the moment we omit the usual first viscosity terms in $\mathbf{j}_E$.) For a purely linear phonon dispersion relation the total energy transported by the phonons is simply the drift term $TS_{ph}\mathbf{v}_{ph}$, and $\mathbf{Q}_3$ is the ^{3}He heat current. Similarly the equation for entropy flow is

$$\frac{\partial S}{\partial t} + \boldsymbol{\nabla} \cdot \mathbf{j}_S = -\frac{1}{T} \mathbf{u} \cdot \frac{\partial \mathbf{g}_3}{\partial t}\Big|_{coll} - \mathbf{Q} \cdot \frac{\boldsymbol{\nabla} T}{T^2} \equiv \mathcal{R}, \tag{19}$$

where

$$\mathbf{u} \equiv \mathbf{v}_3 - \mathbf{v}_{ph}; \tag{20}$$

here the entropy current is $\mathbf{j}_S = \mathbf{v}_{ph} S_{ph} + \mathbf{v}_3 S_3 + \mathbf{Q}_3 / T$. The quantity $\mathcal{R}$ is the rate of entropy generation, which must be non-negative.

In general the change of ^{3}He momentum density is driven by gradients in the chemical potential difference μ and the temperature [19], as we see by subtracting $y \equiv m^* n_3 / (\rho_{ph} + m^* n_3)$ times Eq. (12) from $(1-y)$ times Eq. (10):

$$(1-y) n_3 \left[m^* \frac{\partial \mathbf{u}}{\partial t} + m_3 \boldsymbol{\nabla} \mu \right] - \sigma \boldsymbol{\nabla} T = \frac{\partial \mathbf{g}_3}{\partial t}\Big|_{coll} - y\mathbf{H}. \tag{21}$$

Here

$$\mu = \frac{\mu_3}{m_3} - \frac{\mu_4}{m_4}. \tag{22}$$

and

$$\sigma = y S_{ph} - (1-y) S_3. \tag{23}$$

Relative motion of the ^{3}He and phonons can in general generate a ^{3}He heat current, a type of "thermoelectric" effect; thus as a function of $\boldsymbol{\nabla} T$ and $\mathbf{u}$, the ^{3}He heat current to linear order has the form:

$$\mathbf{Q}_3 = -K_3 \boldsymbol{\nabla} T + T \chi \mathbf{u}. \tag{24}$$

In addition, the rate of momentum transfer in ^{3}He-phonon collisions has the form:

$$\frac{\partial \mathbf{g}_3}{\partial t}\Big|_{coll} = -\frac{m^* n_3}{\tau} \mathbf{u} - \chi \boldsymbol{\nabla} T, \tag{25}$$

where the lifetime τ determines the diffusion constant,

$$D = T\tau/m^* \tag{26}$$

for ^{3}He against phonons. That the same off-diagonal thermoelectric coefficient χ appears in both $\mathbf{Q}$ and $(\partial \mathbf{g}_3/\partial T)_{coll}$ follows from general Onsager reciprocity relations for transport coefficients [20]. As we shall see, χ vanishes for dilute solutions at very low temperature; the first non-vanishing result for χ appears when we take into account phonon recoil corrections (Sec. 6).

With Eqs. (24) and (25), together with (21) in the static limit, we find the rate of entropy generation,

$$\mathcal{R} = \frac{n_3}{D}u^2 + K_3\left(\frac{\boldsymbol{\nabla} T}{T}\right)^2, \tag{27}$$

plus the usual first viscosity term. The kinetic coefficients D, K_3, χ and η_{ph}, specify the transport properties of the solutions; the task we pursue in the following sections is to calculate these coefficients in terms of microscopic scattering processes.

Since in a steady state, $n_3\boldsymbol{\nabla}\mu_3 + S_3\boldsymbol{\nabla}T = (\partial g_3/\partial T)_{coll}$, we find from Eq. (25),

$$\mathbf{u} = -\frac{D}{T}\left(\boldsymbol{\nabla}\mu_3 + \frac{S_3+\chi}{n_3}\boldsymbol{\nabla}T\right). \tag{28}$$

It will be more useful in later calculations of the transport properties to work in terms of the ^{3}He density, n_3, rather than μ_3. Using $n_3 d\mu_3 = d(n_3 T) - S_3 dT + n_3 d\epsilon_0$, we equivalently have

$$\mathbf{u} = -D\left(\frac{\boldsymbol{\nabla} n_3}{n_3} + \frac{\boldsymbol{\nabla}\epsilon_0}{T}\right) - D_T\frac{\boldsymbol{\nabla}T}{n_3}, \tag{29}$$

where D_T, an effective thermoelectric diffusion constant, is given by

$$D_T = \frac{D}{T}(n_3 + \chi). \tag{30}$$

In the situation in which there is no net ^{3}He or ^{4}He particle flow, the total energy current, from Eq. (18), is

$$\mathbf{Q} = \mathbf{Q}_3 - TS_{ph}\mathbf{u} = -K_3\boldsymbol{\nabla}T - T(S_{ph}-\chi)\mathbf{u}; \tag{31}$$

the $TS_{ph}\mathbf{u}$ term is the heat carried by phonons with respect to the ^{3}He. We define the total thermal conductivity, K, of the solutions by calculating, in the absence of ^{3}He and ^{4}He particle transport, the total energy flow, $\mathbf{Q} = \mathbf{Q}_3 - TS_{ph}\mathbf{u} \equiv -K\boldsymbol{\nabla}T$. In the absence of phonon viscous effects (which we include below), the Gibbs-Duhem relation implies in this situation that $n_3\boldsymbol{\nabla}\mu_3 = -S\boldsymbol{\nabla}T$, so that in the static limit Eq. (28) yields

$$\mathbf{u} = \frac{D}{n_3 T}(S_{ph}-\chi)\boldsymbol{\nabla}T, \tag{32}$$

and thus

$$K = K_3 + \frac{D}{n_3}\left(S_{ph} - \chi\right)^2. \tag{33}$$

At constant pressure, the total entropy generation rate, $\mathcal{R}$, is then $K\left(\boldsymbol{\nabla} T/T\right)^2$.

2.3 Phonon viscosity

We now extend the previous discussion of the thermal conductivity to include the phonon viscosity. We consider a steady state in which $\mathbf{v}_3 = 0$, and assume that $\mathbf{v}_{ph}$ is in the z-direction, but varying sinusoidally in the x-direction, i.e., $\mathbf{v}_{ph} \sim e^{ik_x x}\hat{z}v_{ph}$. Equation (12) for conservation of momentum thus reads

$$S_{ph}\boldsymbol{\nabla} T = -\frac{\partial \mathbf{g}_3}{\partial t}\Big|_{coll} - \eta_{ph}k_x^2\mathbf{v}_{ph}. \tag{34}$$

Using Eq. (25), we find

$$\mathbf{v}_{ph} = -\frac{S_{ph} - \chi}{(Tn_3/D) + \eta_{ph}k_x^2}\boldsymbol{\nabla} T. \tag{35}$$

Including the term $-T\chi\mathbf{v}_{ph}$ in the total heat current $\mathbf{Q}$, we then derive the total thermal conductivity of the solution,

$$K = K_3 + T\frac{(S_{ph} - \chi)^2}{(Tn_3/D) + \eta_{ph}k_x^2} \equiv K_3 + K_{ph}. \tag{36}$$

This equation defines the phonon thermal conductivity, K_{ph}.

Equation (36), which encompasses the range of x_3 from that in the Lamoreux experiment to that in the nEDM experiment, yields a number of physically interesting limits. Since $\eta_{ph} = \frac{1}{5}s\rho_{ph}\ell$, where ℓ is the phonon mean free path for viscosity (see Eq. (105)), the denominator in Eq. (36) shows the competition between the mean free path for phonon scattering against the ^{3}He and the viscous diffusion length $\sim 1/k_x^2\ell$. First, for non-zero n_3, as $k_x \to 0$, corresponding to a container infinite in all directions, we derive Eq. (33). On the other hand, in the absence of ^{3}He,

$$K \to K_{ph} = \frac{TS_{ph}^2}{\eta_{ph}k_x^2} = 5\frac{sS_{ph}}{k_x^2\ell}. \tag{37}$$

This result can be understood in terms of a phonon undergoing a random walk of $\sim (1/k_x\ell)^2$ steps in going a wavelength $\sim 1/k_x$.

For Poiseuille flow along the z-direction between parallel plates separated at $x = \pm L_x/2$, one has $v_{ph} \sim L_x^2 - 4x^2$, so that in terms of $\bar{v}_{ph}$, the average flow velocity, $\nabla^2 v_{ph} = -12\bar{v}_{ph}/L_x^2$. Thus in linear order $K_{ph} = 5sS_{ph}L_x^2/12\ell$, which agrees with Eq. (37) with the identification $k_x^2 = 12/L_x^2$. For Poiseuille flow in a cylinder of radius R one has rather $\nabla^2 v_{ph} = -8\bar{v}_{ph}/R^2$ on average, which leads to the Benin-Maris result [21] $K_{ph} = 5sS_{ph}R^2/8\ell$.

To see the regime in which phonon viscosity is important we note that the ratio of the viscous to diffusive term in the denominator of Eq. (35) is

$$\frac{\eta_{ph}k_x^2 D}{Tn_3} = \frac{\ell k_x^2 D}{5s}\frac{S_{ph}}{n_3}. \tag{38}$$

As extracted from viscosity measurements [22,23], the characteristic phonon mean free path for viscosity in pure ^{4}He is

$$\ell(T) \simeq 3.2\times 10^{-3}/T_K^5\,\mathrm{cm} \tag{39}$$

over a broad range of temperatures around $T = 0.5$ K; here T_K is the temperature measured in Kelvin. With Eq. (84) for D we find

$$\frac{\eta_{ph}k_x^2 D}{Tn_3} \sim 0.5\left(\frac{10^{-6}}{x_3}\right)\left(\frac{0.45K}{T}\right)^9\left(\frac{1\,\mathrm{cm}}{R}\right)^2, \tag{40}$$

indicating that viscosity becomes important for $x_3 \lesssim 10^{-6}$ at temperatures of order 0.5 K. At these low concentrations, effects of the ^{3}He on the phonon viscosity are negligible.

As discussed in Sec. 7, K_3 reaches a maximum fraction of only about 1% of the overall thermal conductivity at the highest ^{3}He concentrations considered here, $x_3 = 10^{-3}$. Although the phonon thermal conductivity K_{ph} falls with increasing x_3 as $1/x_3$ in this regime, K_3 is limited by ^{3}He-^{3}He scattering and is never dominant.

2.4 Currents and distribution functions

At low temperatures, the ^{4}He energy current is given in terms of the phonon distribution function, $n_{\mathbf{q}}$, by

$$\mathbf{j}_{E,4} = \left(n_4 - \frac{\delta m}{m_4}n_3\right)\mu_4\mathbf{v}_s + \int\frac{d^3q}{(2\pi)^3}sq(s\hat{\mathbf{q}})n_{\mathbf{q}}, \tag{41}$$

where the second term is the phonon energy current,

$$\mathbf{j}_{E,ph} = s^2\int\frac{d^3q}{(2\pi)^3}\mathbf{q}\,n_{\mathbf{q}} = TS_{ph}\mathbf{v}_{ph}. \tag{42}$$

The integral is simply the momentum density carried by the phonons. The corresponding ^{3}He energy current is similarly given in terms of the ^{3}He distribution function, $f_{\mathbf{p}}$, by

$$j_{E,3} = \frac{\delta m\, n_3}{m_4}\mu_4\mathbf{v}_3 + \nu\int\frac{d^3p}{(2\pi)^3}\frac{\mathbf{p}}{m^*}\left(\frac{p^2}{2m^*} + \epsilon_0\right)f_{\mathbf{p}}, \tag{43}$$

with the ^{3}He flow velocity, $\mathbf{v}_3$, defined by

$$\nu\int\frac{d^3p}{(2\pi)^3}\frac{\mathbf{p}}{m^*}f_{\mathbf{p}} = n_3\mathbf{v}_3. \tag{44}$$

The ^{3}He heat current is

$$\mathbf{Q}_3 = \nu \int \frac{d^3p}{(2\pi)^3} \frac{\mathbf{p}}{m^*} \left(\frac{p^2}{2m^*} + \epsilon_0 \right) f_{\mathbf{p}} - (\mu_3 n_3 + T S_3)\mathbf{v}_3$$
$$= \nu \int \frac{d^3p}{(2\pi)^3} \frac{\mathbf{p}}{m^*} \left(\frac{p^2}{2m^*} - \frac{5}{2}T \right) f_{\mathbf{p}}, \tag{45}$$

since for classical statistics, $(\mu_3 - \epsilon_0)n_3 + TS_3 = (5/2)n_3 T$.

3 Microscopic Scattering

The transport properties of the dilute solutions are determined microscopically by four scattering processes: ^{3}He-phonon scattering, similar to the scattering of non-relativistic electrons and photons; phonon-phonon scattering, which tends to bring the phonons into equilibrium (although incompletely in the present situation), ^{3}He-^{3}He scattering, and lastly, scattering of phonons from the walls (which we consider in Ref. [12]).

3.1 ^{3}He-^{3}He scattering

The matrix element for ^{3}He-^{3}He scattering for atoms of opposite spin for small momentum transfers is [15]

$$V_0 = -0.064 \frac{m_4 s^2}{n_4}. \tag{46}$$

Thus the ^{3}He-^{3}He scattering length is

$$a = \frac{m^* V_0}{4\pi\hbar^2}, \tag{47}$$

the differential cross section is $d\sigma/d\Omega = a^2$, and the total cross section is

$$\sigma_{33} = 4\pi a^2 = \frac{m^{*2}}{4\pi\hbar^4}|V_0|^2$$
$$= \frac{9\pi^3}{k_D^2}(0.064)^2 \left(\frac{m^*}{m_4}\right)^2 \left(\frac{m_4 s}{\hbar k_D}\right)^4 \simeq 10.5\text{Å}^2. \tag{48}$$

A ^{3}He atom of low momentum scatters only from atoms of opposite spin, so that the mean free path of a ^{3}He through a gas of unpolarized ^{3}He is given by

$$\ell_{33} = \frac{2}{n_3 \sigma_{\rm tot}} = \frac{8\pi n_4}{(0.064)^2 x_3} \left(\frac{m_4}{m^*}\right)^2 \left(\frac{\hbar}{m_4 s}\right)^4 \approx \frac{8.66 \times 10^{-8}}{x_3}\,\text{cm}. \tag{49}$$

3.2 ^{3}He-phonon scattering

For ^{3}He thermal velocities, $\sim \sqrt{3T/m^*}$, small compared to s, the scattering is sufficiently elastic that one can, to a good approximation in calculating the ^{3}He-phonon scattering matrix element, neglect the energy transfer in a collision. As shown in [15] [Eq. (24) there], the effective matrix element for elastic scattering of a phonon from momentum $\mathbf{q}$ to $\mathbf{q}'$ and a ^{3}He from momentum $\mathbf{p}$ to $\mathbf{p}'$ is

$$\langle p'q'|T|pq\rangle \equiv \langle \mathcal{T}\rangle = \frac{s\sqrt{qq'}}{2n_4\Omega}(A + B\cos\theta_{qq'}), \tag{50}$$

where $\theta_{qq'}$ is the angle through which the phonon is scattered, the angle between $\mathbf{q}$ and $\mathbf{q}'$; the coefficients are parameters of a solitary ^{3}He in ^{4}He, deduced from experiment to be $A = n_4 d\alpha/dn_4 = -1.2 \pm 0.2$ [24,25], $B = (1+\alpha+\delta m/m_4)(m_4/m^*)(1+\alpha-m_3/m_4) = 0.70 \pm 0.035$ [24,25,26], and Ω is the volume of the system. The largest uncertainty is in A, owing to a systematic difference between the measurements [24,25] of the pressure dependence of the density of dilute solutions.

The differential rate of scattering of a phonon of momentum q by angle $\theta_{qq'}$ is thus

$$\begin{aligned} n_3\frac{d\gamma_q(\theta)}{d\cos\theta_{qq'}} &= n_3\int \frac{q'^2dq'}{2\pi^2}2\pi\delta(sq-sq')|\langle p'q'|T|pq\rangle|^2 \\ &= \frac{x_3 sq^4}{4\pi n_4}(A+B\zeta)^2, \end{aligned} \tag{51}$$

where $\zeta = \cos\theta_{qq'}$; the transport scattering rate for phonons colliding on ^{3}He atoms is

$$\begin{aligned} \gamma_q &= \int \frac{d^3q'}{(2\pi)^3}2\pi\delta(sq-sq')|\langle p'q'|T|pq\rangle|^2(1-\cos\theta_{qq'}) \\ &= \int_{-1}^{1}\frac{d\zeta}{2}\frac{d\gamma_q(\theta)}{d\zeta}(1-\zeta) = s\frac{q^4J}{4\pi n_4^2}, \end{aligned} \tag{52}$$

where $J = A^2 + (B^2 - 2AB)/3 \simeq 2.2 \pm 0.6$. The momentum dependent mean free path of a phonon scattering against the ^{3}He is

$$\ell_{ph3}(q) = \frac{s}{n_3\gamma_q} = \frac{5.52}{3\pi x_3 k_d}\left(\frac{k_d}{q}\right)^4, \tag{53}$$

where $n_4 = k_D^3/6\pi^2$ and $k_D \simeq 1.089$ Å.

Similarly, as we shall see (Eq. (81) with $n = 0$), the effective ^{3}He-phonon relaxation rate in diffusion is $\Gamma/3m^*T$, where

$$\begin{aligned} \Gamma &= \int \frac{d^3q}{(2\pi)^3}q^2\gamma_q n_q^0(1+n_q^0) \\ &= \frac{9!}{2}\pi\zeta(8)J\left(\frac{T}{\hbar s k_D}\right)^9 k_D^3 s, \end{aligned} \tag{54}$$

with $\zeta(8) = \pi^4/9450 \simeq 1.004$ the Riemann zeta function. The mean free path for scattering of a thermal ^{3}He by phonons [27] is

$$\ell_{3ph} = \frac{2}{\sqrt{3}\pi 8!\zeta(8)Jk_D}\left(\frac{ms}{k_D}\right)^{1/2}\left(\frac{\hbar s k_D}{T}\right)^{15/2} = 0.077\left(\frac{0.45K}{T}\right)^{15/2}\text{cm}. \quad (55)$$

Comparing the mean free paths, Eqs. (49) and (55) to estimate the importance of 3-3 versus phonon scattering in bringing the ^{3}He into equilibrium. we find

$$\frac{\ell_{3ph}}{\ell_{33}} = 0.89\times 10^6 x_3\left(\frac{0.45K}{T}\right)^{15/2} \quad (56)$$

For $T = 0.45$ K and $x_3 = 10^{-6}$, $\ell_{3ph} \approx \ell_{33}$, while for $T = 0.65$ K and $x = 3\times 10^{-4}$, $\ell_{3ph}/\ell_{33} \approx 16.9$.

3.3 Phonon-phonon scattering

Phonon-phonon scattering conserves total phonon momentum, and thus does not contribute to the thermal conductivity directly. However, its effect on the phonon distribution is important. Because of the anomalous dispersion of phonons in liquid ^{4}He at low pressure, three-phonon Beliaev and Landau-damping processes are allowed; these processes rapidly equilibrate phonons propagating in a given direction. producing a phonon distribution,

$$\tilde{n}_{\mathbf{q}} = \frac{1}{e^{\tilde{\beta}(\hat{q},\mathbf{r})sq}-1}, \quad (57)$$

along rays in momentum space [22], in which the temperature is dependent on the momentum direction $\hat{q}$. From energy conservation in these rapid scatterings,

$$\int sq(q^2dq)\,(n_{\mathbf{q}} - \tilde{n}_{\mathbf{q}}) = 0. \quad (58)$$

For phonons with small momentum $q \sim T/s$, the dominant three-phonon process is Landau damping, in which the phonon is absorbed or emitted by a thermal phonon, thereby producing another phonon with energy $\sim T$. The rate of this process is

$$\frac{1}{\tau_L} = \frac{3\pi}{2}(u+1)^2 sq\frac{\rho_n}{m_4 n_4}, \quad (59)$$

where $u = \partial \ln s/\partial \ln n_4 \simeq 2.843$ at SVP is the phonon Grüneisen parameter. The Beliaev process, the decay of one phonon into two, has a rate

$(u+1)^2 q^5/240\pi m_4 n_4$; this process dominates at large q ($\gg T/s$) [28]. The ratio of the rates of phonon scattering on ^{3}He to Landau damping is

$$\frac{15xJ}{(u+1)^2}\left(\frac{sq}{T}\right)^3\frac{m_4 s^2}{T} \simeq 0.5x\left(\frac{sq}{T}\right)^3\frac{m_4 s^2}{T} \ll 1. \tag{60}$$

We will describe such rapid scatterings in the phonon Boltzmann equation in terms of a relaxation time, τ_r, with a collision rate,

$$\left(\frac{\partial n_{\mathbf{q}}}{\partial t}\right)_{rapid} = -\frac{1}{\tau_r}\left(n_{\mathbf{q}} - \tilde{n}_{\mathbf{q}}\right). \tag{61}$$

Relaxation between rays, which conserves phonon momentum and energy, occurs on a longer time scale. We describe such processes, which determine the phonon viscosity, in a relaxation time approximation

$$\left(\frac{\partial n_{\mathbf{q}}}{\partial t}\right)_{long} = -\frac{1}{\tau_l}\left(n_{\mathbf{q}} - n^{le}_{\mathbf{q}}\right). \tag{62}$$

where $\tau_l = \ell(T)/s$. Here

$$n^{le}_{\mathbf{q}} = \frac{1}{e^{\beta(\mathbf{r})(sq - \mathbf{q}\cdot\mathbf{v}_{ph})} - 1} \tag{63}$$

is the local equilibrium phonon distribution function, in terms of the local temperature $T(\mathbf{r}) = 1/\beta(\mathbf{r})$ and mean flow velocity $\mathbf{v}_{ph}$ that the phonons would have by relaxing to local equilibrium through phonon-phonon collisions.

When phonon viscosity is important, the angular dependent temperature of the phonon distribution has the form

$$\delta\beta(\hat{q}) = -\frac{\beta}{s}\left(\hat{q}_z v_{ph} + \hat{q}_z\hat{q}_x\lambda\right), \tag{64}$$

where λ measures the quadrupolar distortion of the phonon distribution, proportional to the gradient in the transverse direction of the phonon velocity v_{ph}, which we take along the x direction to be specific. The relaxation time, τ_ℓ depends strongly on the angular dependence of $n_{\mathbf{q}}$, which is a second spherical harmonic (the final term in Eq. 64)) for phonon viscosity. For $\ell \gg 1$ one expects $1/\tau_\ell \sim \ell^2$, until $1/\ell \sim \theta$, where θ is a typical scattering angle.

4 ^{3}He Boltzmann Equation

We turn now to calculating the transport coefficients, D, K_3, and χ from the ^{3}He Boltzmann equation. We assume quite generally that the driving terms,

$\boldsymbol{\nabla}\mu_3$ and $\boldsymbol{\nabla}T$ and thus $\mathbf{v}_{ph}$ are in the z direction. The ^{3}He Boltzmann equation has the form

$$\frac{\partial f_{\mathbf{p}}}{\partial t} + \frac{\mathbf{p}}{m^*}\cdot\boldsymbol{\nabla}_r f_{\mathbf{p}} = -\frac{f_{\mathbf{p}} - f^{le}_{\mathbf{p}}}{\tau_{33}} + \sum_{p',q,q'} |\langle\mathcal{T}\rangle|^2 2\pi\delta(\Delta E)\delta_{\mathbf{p}+\mathbf{q},\mathbf{p}'+\mathbf{q}'} \times \left[f_{\mathbf{p}'} n_{\mathbf{q}'}(\mathbf{r})(1+n_{\mathbf{q}}(\mathbf{r})) - f_{\mathbf{p}}\, n_{\mathbf{q}}(\mathbf{r})(1+n_{\mathbf{q}'}(\mathbf{r})\right], \tag{65}$$

where τ_{33} is the ^{3}He-^{3}He scattering time (appropriate for vector drivers of the ^{3}He away from equilibrium); we take a mean thermal velocity to define $\tau_{33} = \sqrt{m^*/3T}\ell_{33}$ in terms of the ^{3}He-^{3}He mean free path. Also

$$f^{le}_{\mathbf{p}} = e^{-\beta(\mathbf{r})(p^2/2m^* - \mathbf{p}\cdot\mathbf{v}_3 - \mu_3(\mathbf{r}))} \tag{66}$$

is the distribution function towards which ^{3}He-^{3}He collisions drive the ^{3}He, and $\langle\mathcal{T}\rangle \equiv \langle p'q'|T|pq\rangle$. In calculating D, K_3, and χ, phonon viscosity can be neglected. Then, as argued above, rapid phonon equilibration along rays in momentum space brings the phonon distribution into the form (63). We first linearize the phonon-^{3}He-phonon collision term in terms of deviations from the phonon and ^{3}He distribution functions from equilibrium:

$$\delta\tilde{n}_{\mathbf{q}} = \tilde{n}_{\mathbf{q}} - n^{le0}_q = \beta n^0_q(1+n^0_q) sq\mathbf{q}\cdot\mathbf{v}_{ph}, \tag{67}$$

and

$$\delta f_{\mathbf{p}} = f_{\mathbf{p}} - f^{le0}_p = \beta f^0_p p_z w_p; \tag{68}$$

here the local equilibrium ^{3}He distribution function is

$$f^{le0}_p = e^{-\beta(\mathbf{r})(p^2/2m^* - \bar{\mu}_3(\mathbf{r}))} \tag{69}$$

and the global equilibrium distribution function is

$$f^0_p = e^{-\beta(p^2/2m^* - \bar{\mu}_3)}, \tag{70}$$

where $\bar{\mu}_3 = \mu_3 - \epsilon_0$. With the detailed balance condition, $f^0_{p'} n^0_{q'}(1+n^0_q) = f^0_p n^0_q(1+n^0_{q'})$, the term in square brackets in Eq. (65) becomes in linear order,

$$-f^0_p n^0_q(1+n^0_{q'})\left[\beta(w_p p_z - w_{p'}p'_z + (\mathbf{q}-\mathbf{q}')\cdot\mathbf{v}_{ph}\right]. \tag{71}$$

It is most convenient to use the momentum conservation to eliminate $\mathbf{p}'$, and write $\mathbf{p}' = \mathbf{p} + \mathbf{k}$, where $\mathbf{k} = \mathbf{q} - \mathbf{q}'$. Then energy conservation implies that

$$\Delta E \equiv sq' - sq + \frac{\mathbf{p}\cdot\mathbf{k}}{2m^*} + \frac{k^2}{2m^*} = 0. \tag{72}$$

The momentum transfer k is characteristically of order T/s, which is small compared with the momentum of a ^{3}He atom, $\sim (m^*T)^{1/2}$. The approach we

take will be to expand the collision integrals in powers of $k/(m^*T)^{1/2}$; the leading terms are of second order. One might have imagined that to leading order one could neglect the energy transfer in collisions. However, this is not the case because, in a collision, the relative changes in the momentum and energy of a ^{3}He atom are both of order $(T/m^*s^2)^{1/2}$. As we show, the differential equation we derive for the distribution function to order k^2/m^*T, in the form of a Fokker-Planck equation, has an exact analytic solution.

Our calculations show that the energy transfer in a collision of ^{3}He with a phonon, although relatively small, has a large qualitative effect on the ^{3}He distribution function. In Ref. [6] we showed that, if the energy transfer were neglected, the relaxation time for a quasiparticle varied as p^2, whereas the exact solution above shows that when the energy transfer is taken into account, the relaxation time is independent of p.

The next higher-order terms vary as k^4 and we shall refer to them as "recoil corrections". These give rise to contributions of order T/m^*s^2 times the leading term. Despite the fact that T/m^*s^2 is $\sim 10^{-2}$ at temperatures of order 0.5 K, these corrections are not negligible because of the large numerical coefficients, as we demonstrated in the Appendix of Ref. [6]. We return to the recoil corrections in Sec. 6.

Expanding $w_p p_z - w_{p'} p'_z$ to order k^2 we find

$$\begin{aligned} w_p p_z - w_{p'} p'_z &= -w_p k_z - (p_z + k_z)\mathbf{k}\cdot\boldsymbol{\nabla}_p w_p - \frac{1}{2} p_z (\mathbf{k}\cdot\boldsymbol{\nabla}_p)^2 w_p \\ &= -w_p k_z - (p_z + k_z)\mathbf{k}\cdot\hat{\mathbf{p}}\, w'_p - \frac{1}{2} p_z \left[(\mathbf{k}\cdot\hat{\mathbf{p}})^2 (w''_p - w'_p/p) + k^2 w'_p/p\right], \end{aligned} \tag{73}$$

where primes on the w's denote derivatives with respect to p.

The ^{3}He-phonon collision term in Eq. (65), with the expansion (73), becomes

$$\left(\frac{\partial f_p}{\partial t}\right)_{3-ph\, coll} = \beta \sum_{\mathbf{q},\mathbf{q}'} |\langle\mathcal{T}\rangle|^2 2\pi\delta(\Delta E)\delta_{\mathbf{p}+\mathbf{q},\mathbf{p}'+\mathbf{q}'} f_p^0\, n_q^0 (1 + n_{q'}^0)$$
$$\times \left((w_p - v_{ph})k_z + (p_z + k_z)\mathbf{k}\cdot\hat{\mathbf{p}}\, w'_p + \frac{1}{2} p_z \left[(\mathbf{k}\cdot\hat{\mathbf{p}})^2 (w''_p - w'_p/p) + k^2 w'_p/p\right]\right). \tag{74}$$

The terms in the final parentheses are both linear and quadratic in $\mathbf{k}$. We expand the energy delta function to first order in $\mathbf{p}\cdot\mathbf{k}/m^*$, and average over the direction of $\mathbf{k}$, keeping the direction of $\mathbf{p}$ fixed. The terms in the final parentheses of order k^2 give

$$\frac{k^2 p_z}{6}\left(4w'_p/p + w''_p\right) n_q^0 (1 + n_q^0)\delta(sq' - sq). \tag{75}$$

We symmetrize the terms of order $\mathbf{k}$ under the transformation $\mathbf{q} \leftrightarrow \mathbf{q}'$; the argument of the delta function remains fixed, while $\mathbf{k} \to -\mathbf{k}$ and $n_q^0(1+n_{q'}^0) \to$

$n^0_{q'}(1+n^0_q)$. Thus the terms of order $\mathbf{k}$ lead to

$$\frac{k^2 p_z}{6m^*}\left(w_p - v_{ph} + pw'_p\right)\left(n^0_q - n^0_{q'}\right)\delta'(sq'-sq). \tag{76}$$

When we integrate by parts in q' this term becomes

$$-\beta\frac{k^2 p_z}{6m^*}\left(w_p - v_{ph} + pw'_p\right) n^0_q(1+n^0_q)\delta(sq'-sq). \tag{77}$$

Altogether then, in terms of Γ defined in Eq. (54), we have

$$\left(\frac{\partial f_p}{\partial t}\right)_{3ph\,coll} = \frac{\beta\Gamma}{3}p_z f^0_p\left(w''_p + 4\frac{w'_p}{p} - \frac{\beta}{m^*}\left(pw'_p + w_p - v_{ph}\right)\right). \tag{78}$$

Remarkably, the derivatives of the distribution function in the parentheses on the right are just those entering into the equation for the polynomials, $g_n(y)$ [with $g_0 \sim 1$, $g_1 \sim y^2 - 5$, etc.], which obey

$$g''_n + (4/y - y)g'_n + 2ng_n = 0. \tag{79}$$

These polynomials are related to the Sonine polynomials familiar from the theory of dilute gases by the relation $g_n(y) \propto S^n_{3/2}(y^2/2)$ [9,11]. Writing $w_n(p) = g_n(p\sqrt{\beta/m^*})$ we see that

$$w''_n + 4w'_n/p - (\beta/m)pw'_n = -2n(\beta/m^*)w_n. \tag{80}$$

so that the eigenfunctions of the differential operator in Eq. (81) are just the polynomials $g_n(y)$. The collision term for a deviation proportional to $g_n(y)$ is thus

$$\begin{aligned}\left(\frac{\partial f_p}{\partial t}\right)_{3ph\,coll} &= -\frac{\beta^2\Gamma}{3m^*}p_z f^0_p\left[(2n+1)w_n - v_{ph}\right] \\ &= -\frac{\beta\Gamma}{3m^*}[(2n+1)\delta f_n - \beta f^0_p p_z v_{ph}].\end{aligned} \tag{81}$$

We first consider steady state diffusion driven by a ^{3}He chemical potential gradient at constant temperature. The driving term on the left side of the Boltzmann equation is $(p_z/m^*)\beta f^0_p \partial\bar{\mu}_3/\partial z$, and the solution is the first Sonine polynomial, a constant. In other words, the ^{3}He distribution is just an equilibrium drifting at velocity v_3, for which the ^{3}He-^{3}He collision term vanishes. Then

$$\frac{p_z}{m^*}\beta f^0_p\frac{\partial\bar{\mu}_3}{\partial z} = -\frac{\beta^2\Gamma}{3m^*}p_z f^0_p u, \tag{82}$$

where $u = v_3 - v_{ph}$, so that $u = -(3T/\Gamma)\partial\bar{\mu}_3/\partial z$; the diffusion constant is thus

$$D = \frac{3T^2}{\Gamma}, \tag{83}$$

and the collision time in Eq. (25) is $\tau = 3m^*T/\Gamma$. Using Eq. (54), with $\hbar s k_D = 19.9\text{K}$ and $\hbar k_D/m_4 s = 0.729$ we find numerically that

$$D = 173\left(\frac{0.45K}{T_K}\right)^7 \text{cm}^2/\text{sec} = \frac{0.65}{T_K^7}\text{cm}^2/\text{sec}. \tag{84}$$

We note that there is a $\sim 35\%$ increase in this lowest order calculation of D due to the recoil corrections (see Secs. 6 and 7).

We next compute the thermal conductivity of the ^{3}He, defined by the ^{3}He heat current produced by a temperature gradient at constant ^{3}He pressure, cf. Eq. (45). At constant P_3, the driving term on the left side of the Boltzmann equation is proportional to

$$\left(\frac{\partial f_p^{le}}{\partial z}\right)_{P_3} = -f_p^0\left(\frac{p^2}{2m^*} - \frac{5}{2}T\right)\frac{\partial \beta}{\partial z}. \tag{85}$$

Thus the deviation of the ^{3}He distribution is a first ($n = 1$) Sonine polynomial $\sim p^2 - 5m^*T$; we write

$$w_p = C\left(\frac{p^2}{2m^*} - \frac{5}{2}T\right). \tag{86}$$

Wiith Eq. (81) the Boltzmann equation yields

$$C = -\frac{1}{\Gamma'}\frac{\partial T}{\partial z}, \tag{87}$$

where $\Gamma' = \Gamma + m^*/\beta\tau_{33}$. Thus the ^{3}He heat current is

$$Q_3 = -\nu\int \frac{d^3p}{(2\pi)^3}\frac{p_z^2}{m^*}\left(\frac{p^2}{2m} - \frac{5}{2}T\right)^2\frac{\beta}{\Gamma'}\frac{\partial T}{\partial z}. \tag{88}$$

Evaluating the integral we find the ^{3}He thermal conductivity,

$$K_3 = \frac{\beta}{\Gamma'}\nu\int \frac{d^3p}{(2\pi)^3}\frac{p^2}{2m}\left(\frac{p^2}{2m} - \frac{5}{2}T\right)^2 = \frac{5T^2}{2\Gamma'}n_3, \tag{89}$$

Numerically, the ^{3}He-^{3}He scattering contribution to Γ' is of relative order $(10^{-6}/x_3)(T/0.45K)^{15/2}$, indicating that ^{3}He-^{3}He scattering does not contribute importantly at concentrations well below 10^{-6} at temperatures of order 0.5 K.

At the present level of approximation the thermoelectric coefficient χ is zero. To see this we assume a uniform temperature; then the solution of the Boltzmann equation is simply a drifting local equilibrium [cf. Eq. (82)] for which, from the orthogonality of the Sonine polynomials, the ^{3}He heat current vanishes. Thus from Eq. (24), $\chi = 0$. Similarly $(\partial g_3/\partial t)_{coll}$ also vanishes if the distribution function is proportional to the first Sonine polynomial, and thus from Eq. (25), we see again that χ vanishes. However, with terms of higher order in k included in the collision term, the distribution functions are not

given simply in terms of Sonine polynomials, and one finds $\chi \neq 0$ (see Sec. 6.2. We expect, from expanding the ^{3}He-phonon collision term to higher order in k that the corrections are $\sim k^2/p^2 \sim T/m^*s^2$. Thus in order of magnitude $\chi \sim n_3 T/m^* s^2$, which leads to negligible corrections to D_T, Eq. (30), as well as K_{ph} in Eq. (33).

5 Phonon Boltzmann Equation

We turn to the details of the phonon Boltzmann equation. With phonon-phonon scatterings treated in a relaxation time approximation, the phonon Boltzmann equation in dilute solutions has the structure:

$$\frac{\partial n_{\mathbf{q}}}{\partial t} + s\hat{q}\cdot\boldsymbol{\nabla}_r n_{\mathbf{q}} = \left(\frac{\partial n}{\partial t}\right)_{\mathrm{ph}-^3\mathrm{He}} - \frac{1}{\tau_r}\left(n_{\mathbf{q}} - \tilde{n}_{\mathbf{q}}\right) - \frac{1}{\tau_l}(n_{\mathbf{q}} - n_q^{le}), \tag{90}$$

where the phonon-^{3}He collision term is, cf. Eq, (65),

$$\left(\frac{\partial n}{\partial t}\right)_{\mathrm{ph}-^3\mathrm{He}} = \sum_{p,p',q'} |\langle\mathcal{T}\rangle|^2 2\pi\delta(\Delta E)\delta_{\mathbf{p}+\mathbf{q},\mathbf{p}'+\mathbf{q}'} \times \left[f_{\mathbf{p}'} n_{\mathbf{q}'}(\mathbf{r})(1+n_{\mathbf{q}}(\mathbf{r})) - f_{\mathbf{p}} n_{\mathbf{q}}(\mathbf{r})(1+n_{\mathbf{q}'}(\mathbf{r})\right]. \tag{91}$$

We do not include explicit scattering of phonons with the container walls in this paper.

In the limit that the rate of phonon collisions with the ^{3}He is much smaller than phonon-phonon collisions along rays, cf. Eq. (60), and the spatial and temporal variations are slow, the deviation of the distribution function $n_{\mathbf{q}}$ from $\tilde{n}_{\mathbf{q}}$ is of relative order τ_r; as we see from Eq. (90),

$$n_{\mathbf{q}} - \tilde{n}_{\mathbf{q}} = \tau_r\left[\frac{\partial n_{\mathbf{q}}}{\partial t} + s\hat{q}\cdot\boldsymbol{\nabla}_r n_{\mathbf{q}} + \frac{1}{\tau_l}(n_{\mathbf{q}} - n_q^{le}) - \left(\frac{\partial n}{\partial t}\right)_{\mathrm{ph}-^3\mathrm{He}}\right]. \tag{92}$$

Thus all n on the right can be replaced by $\tilde{n}$. The condition (58) then allows us to derive the effective Boltzmann equation for $\beta(\hat{q},\mathbf{r})$, since the integral $\int dq\, q^3$ of the right side of Eq. (92) vanishes. We write

$$\delta\tilde{n}_{\mathbf{q}} = \tilde{n}_{\mathbf{q}} - n_q^{le0} = -n_q^0(1+n_q^0)sq\,\delta\beta(\hat{q},\mathbf{r}), \tag{93}$$

where $\delta\beta(\hat{q},\mathbf{r}) = \beta(\hat{q},\mathbf{r}) - \beta(\mathbf{r})$, with $\beta(\mathbf{r})$ the local equilibrium temperature, and using $\int q^3 dq \delta\tilde{n}_{\mathbf{q}} = -6\pi^2 s\rho_{ph}\delta\beta$. we arrive at the equation for $\beta(\hat{q},\mathbf{r})$:

$$\frac{\partial\beta(\hat{q},\mathbf{r})}{\partial t} + s\hat{q}\cdot\boldsymbol{\nabla}_r\beta(\hat{q},\mathbf{r}) = \mathcal{C} - \frac{1}{\tau_l}(\beta(\hat{q},\mathbf{r}) + \beta q_z v_{ph}/s). \tag{94}$$

Here

$$\mathcal{C} = -\int \frac{q^3 dq}{6\pi^2 s\rho_{ph}} \left(\frac{\partial \tilde{n}}{\partial t}\right)_{\mathrm{ph}-^3\mathrm{He}}; \tag{95}$$

the tilde on the right side indicates that the phonon distribution functions $n_{\mathbf{q}}$ in the collision terms are replaced by $\tilde{n}_{\mathbf{q}}$.

The analysis of the ^{3}He-phonon collision term in the phonon Boltzmann equation is similar to that in the ^{3}He Boltzmann equation. We assume $\beta(\hat{q})$ to have the form (64), and treat the v_{ph} and λ terms separately. In the v_{ph} term, using Eq. (73), we see that when one averages over the direction of ^{3}He momenta $\mathbf{p}$ only the terms first order in $\mathbf{k}$ survive; then

$$\begin{aligned}\left(\frac{\partial n}{\partial t}\right)_{\mathrm{ph}-^3\mathrm{He}} &= \sum_q k_z |\langle \mathcal{T}\rangle|^2 2\pi\delta(sq - sq')n_q^0(1+n_q^0))F_3 \\ &= q\gamma_q n_q^0(1+n_q^0))F_3,\end{aligned} \tag{96}$$

where

$$F_3 = \sum_p f_p^0 \left(w_p - v_{ph} + \frac{1}{3}pw_p'\right). \tag{97}$$

If the ^{3}He are driven by ^{3}He chemical potential gradients, then $F_3 = n_3 u$, while if the ^{3}He are driven by a temperature gradient, then $w_p = -(1/\Gamma)(\partial T/\partial z)\times \big(p^2/2m^* - 5T/2\big)$, and $F_3 = -n_3 v_{ph}$.

The λ terms in Eq. (64) lead to a similar term in the collision integral,

$$-\gamma_q^{(2)}\left(\delta\tilde{n}_{\mathbf{q}} - n_q^0(1+n_q^0)\beta q_z v_{ph}\right), \tag{98}$$

where

$$\gamma_q^{(2)} = \int \frac{d\cos\theta}{2}(1 - P_2(\cos\theta))\frac{d\gamma_q(\theta)}{d\cos\theta} \tag{99}$$

is the average of the scattering rate of phonons on ^{3}He appropriate for viscosity, with P_2 the second Legendre polynomial. Thus

$$\mathcal{C} = \frac{n_3\beta^2\hat{q}_z}{3s\rho_{ph}}\left(-\Gamma u + \Gamma^{(2)}\hat{q}_x\lambda\right), \tag{100}$$

where Γ is given by Eq. (54), and

$$\Gamma^{(2)} = \int \frac{d^3q}{(2\pi)^3} q^2 \frac{\gamma^{(2)}(q)}{n_3} n_q^0(1+n_q^0) = \frac{A^2 + B^2/5}{A^2 + (B^2 - 2AB)/3}\Gamma; \tag{101}$$

the numerical coefficient is $\simeq 0.74$.

The linearized phonon Boltzmann equation, Eq. (94), separates into equations for the individual spherical harmonic components of $\beta(\hat{q})$. We first calculate the phonon viscosity the second spherical harmonic term; to do so we assume static flow and compute the off-diagonal component $T_{xz} = -\eta_{ph}\partial v_{ph}/\partial x$. Then

$$\lambda = -s\tau_v \boldsymbol{\nabla}_x v_{ph} \tag{102}$$

where

$$\frac{1}{\tau_v} = \frac{n_3\beta}{3\rho_{ph}}\Gamma^{(2)} + \frac{1}{\tau_l}. \tag{103}$$

With Eq. (102), we find

$$\begin{aligned} T_{zx} &= -\int \frac{d^3q}{(2\pi)^3} s q_z \hat{q}_x \delta n_{\mathbf{q}} \\ &= \int \frac{d^3q}{(2\pi)^3} \beta s^2 \tau_v \hat{q}_z^2 \hat{q}_x^2 q^2 n_q^0 (1+n_q^0) \frac{\partial v_{ph}}{\partial z}, \end{aligned} \tag{104}$$

and thus

$$\eta_{ph} = \frac{1}{5}\rho_{ph} s^2 \tau_v. \tag{105}$$

The contribution to the viscosity from phonon scattering on the ^{3}He is important for $(n_3/S_{ph})(s\ell/D) \simeq 3\times 10^4 x_3/T_K \gtrsim 1$.

In extracting the first spherical harmonic component of Eq. (94), we encounter a term $\hat{q}_z\hat{q}_x\hat{q}\cdot\boldsymbol{\nabla}_r\lambda$, which from Eq. (102) equals $-s\tau_v\hat{q}_z\hat{q}_x^2\boldsymbol{\nabla}_x^2 v_{ph}$. The first spherical harmonic component of $\hat{q}_z\hat{q}_x^2$ is $3\int d\Omega(\mathbf{q})\hat{q}_z^2\hat{q}_x^2/4\pi = 1/5$. Thus the phonon momentum density obeys

$$\rho_{ph}\frac{\partial v_{ph}}{\partial t} + S_{ph}\frac{\partial T}{\partial z} - \rho_{ph}\frac{s^2\tau_v}{5}\boldsymbol{\nabla}_x^2 v_{ph} = \frac{n_3\beta}{3}\Gamma u. \tag{106}$$

which we recognize as the phonon momentum conservation equation (12).

6 Recoil corrections

As we showed above, to leading order in an expansion of the ^{3}He-phonon collision operator in powers of $k^2/m^*T \sim T/m^*s^2$, the eigenfunctions of the ^{3}He-phonon collision operator in the ^{3}He Boltzmann equation have the form of Sonine polynomials times p_i for driving terms proportional to $p_i\partial f^{le}_{\mathbf{p}}/\partial x_i$. Terms of order $k^4/m^{*2}T^2$ in the expansion of the collision operator lead to contributions to the transport coefficients of nominal order T/m^*s^2 relative to the leading term. The task in this section is to calculate these corrections explicitly. We have seen that in a temperature or ^{3}He chemical potential gradient, the phonon distribution function is one of drifting local equilibrium, and therefore, it is convenient to work in the reference frame in which the phonon

fluid is at rest ($\mathbf{v}_{ph} = 0$). We also assume that the system is in a steady state and, on the left hand of the Boltzmann equation (65), we may replace the distribution function by a local equilibrium one. At the low concentrations of interest, the ^{3}He-^{3}He scattering may be neglected.

For temperature and chemical potential gradients in the z direction, the left side of the Boltzmann equation (65) becomes

$$\begin{aligned}\frac{\mathbf{p}}{m^*}\cdot\boldsymbol{\nabla}_r f_{\mathbf{p}} &= -\frac{f^0_{\mathbf{p}}}{T}\frac{p_z}{m^*}\left(\frac{\partial(\mu_3-\epsilon_0)}{\partial z}+\frac{p^2/2m^*+\epsilon_0-\mu_3}{T}\frac{\partial T}{\partial z}\right)\\ &= -\frac{f^0_{\mathbf{p}}}{T}\frac{p_z}{m^*}\left(\frac{1}{n_3}\frac{\partial P_3}{\partial z}-\frac{\partial\epsilon_0}{\partial z}+\frac{(p^2/2m^*-5T/2)}{T}\frac{\partial T}{\partial z}\right). \quad (107)\end{aligned}$$

The latter form, in which we use the ^{3}He pressure and the temperature as independent variables, is particularly convenient because it shows that the driving term due to the pressure gradient is proportional to the particle current, p_z/m^*, carried by an atom, and that the driver due to the temperature gradient is proportional to the heat current, $(p^2/2m^*-5T/2)p_z/m^*$. Since the Boltzmann equation is linear, we write $w(p) = W^c(p)\,(1/n_3\partial P_3/\partial z - \partial\epsilon_0/\partial z) + W^h(p)\partial T/\partial z$, and have the equations for the W's:

$$\frac{p_z}{m^*} = \sum_{\mathbf{p}',\mathbf{q},\mathbf{q}'}|\langle\mathcal{T}\rangle|^2 2\pi\delta(\Delta E)\delta_{\mathbf{p}+\mathbf{q},\mathbf{p}'+\mathbf{q}'}n^0_{\mathbf{q}}(1+n^0_{\mathbf{q}'})[p_z W^c(p)-p'_z W^c(p')]. \quad (108)$$

and

$$\left(\frac{p^2}{2m^*}-\frac{5}{2}T\right)\frac{p_z}{m^*} = \sum_{\mathbf{p}',\mathbf{q},\mathbf{q}'}|\langle\mathcal{T}\rangle|^2 2\pi\delta(\Delta E)\delta_{\mathbf{p}+\mathbf{q},\mathbf{p}'+\mathbf{q}'}n^0_{\mathbf{q}}(1+n^0_{\mathbf{q}'}) \times[p_z W^h(p)-p'_z W^h(p')]. \quad (109)$$

The response of the ^{3}He particle number current density and heat current density to the chemical potential and temperature gradients, may be written in terms of the correlation functions $\Xi_{\eta\lambda}$ between the currents as

$$\mathbf{j}_3 = -\Xi_{cc}\left(\frac{1}{n_3}\boldsymbol{\nabla}P_3-\boldsymbol{\nabla}\epsilon_0\right)-\Xi_{ch}\frac{\boldsymbol{\nabla}T}{T}, \quad (110)$$

and

$$\mathbf{Q}_3 = -\Xi_{hc}\left(\frac{\boldsymbol{\nabla}P_3}{n_3}-\boldsymbol{\nabla}\epsilon_0\right)-\Xi_{hh}\frac{\boldsymbol{\nabla}T}{T}, \quad (111)$$

where c denotes the current of particle number and h the heat current. The Onsager reciprocal relations imply that $\Xi_{ch} = \Xi_{hc}$, a fact that can be confirmed explicitly using the results for the $\Xi_{\eta\lambda}$ given below.

On eliminating $(\boldsymbol{\nabla}P_3)/n_3 - \boldsymbol{\nabla}\epsilon_0$ from Eqs. (110) and (111), we find

$$\mathbf{Q}_3 = \frac{\Xi_{hc}}{\Xi_{cc}}n_3\mathbf{u}-\left(\Xi_{hh}-\frac{\Xi^2_{hc}}{\Xi_{cc}}\right)\frac{\boldsymbol{\nabla}T}{T}, \quad (112)$$

a result valid in an arbitrary frame. Comparison of this result with Eq. (24) shows that

$$\chi = \frac{n_3}{T}\frac{\Xi_{hc}}{\Xi_{cc}} \tag{113}$$

and the ^{3}He thermal conductivity is $K_3 = \Xi_{hh} - \Xi_{hc}^2/\Xi_{cc}$.

Microscopically, the number and heat current densities are given by

$$(j_3)_z = \nu \int \frac{d^3p}{(2\pi)^3}\frac{p_z}{m^*}\frac{f^0_{\mathbf{p}}}{T}p_z w(p), \tag{114}$$

and

$$Q_{3z} = \nu \int \frac{d^3p}{(2\pi)^3}\left(\frac{p^2}{2m^*} - \frac{5}{2}T\right)\frac{p_z}{m^*}\frac{f^0_{\mathbf{p}}}{T}p_z w(p), \tag{115}$$

and therefore

$$\Xi_{\eta,\lambda} = \nu \int \frac{d^3p}{(2\pi)^3}\frac{f^0_{\mathbf{p}}}{T}X^\eta(p)\Phi^\lambda(p), \tag{116}$$

where $X^c(p) = p_z/m^*$, $X^h(p) = (p^2/2m^* - 5T/2)p_z/m^*$, and $\Phi^\lambda(p) = W^\lambda p_z$.

We may write Eqs. (108) and (109) in a compact matrix notation as

$$|X^\lambda\rangle = I|\Phi^\lambda\rangle, \tag{117}$$

in terms of which

$$\Xi_{\eta,\lambda} = \langle X^\eta|\Phi^\lambda\rangle, \tag{118}$$

where the inner product is defined by

$$\langle C|D\rangle = \nu \int \frac{d^3p}{(2\pi)^3}\frac{f^0_{\mathbf{p}}}{T}C^\eta(p)D^\lambda(p). \tag{119}$$

We now expand the collision integral in powers of k^2/m^*T,

$$I = I_0 + I_1 + \ldots, \tag{120}$$

where I_0 is the leading term, which is $\sim T/m^*s^2$, and I_1 is the term of order $(k^2/m^*T)^2 \sim (T/m^*s^2)^2$. Similarly, we write the deviation function in the form

$$|\Phi^\lambda\rangle = |\Phi_0^\lambda\rangle + |\Phi_1^\lambda\rangle + \ldots . \tag{121}$$

Equating terms of the same order in T/m^*s^2 in Eq. (117), one finds

$$|X^\lambda\rangle = I_0|\Phi_0^\lambda\rangle, \tag{122}$$

and

$$0 = I_0|\Phi_1^\lambda\rangle + I_1|\Phi_0^\lambda\rangle, \tag{123}$$

from which one sees that

$$|\Phi_1^\lambda\rangle = -(I_0)^{-1} I_1 |\Phi_0^\lambda\rangle \,. \tag{124}$$

The transport coefficient giving the response of a variable specified by $|X^\eta\rangle$ may be expressed in the form

$$\begin{aligned}\Xi^{\eta,\lambda} &= \langle X^\eta|\Phi^\lambda\rangle = \langle X^\eta|\Phi_0^\lambda\rangle + \langle X^\eta|\Phi_1^\lambda\rangle + \dots \\ &= \langle X^\eta|\Phi_0^\lambda\rangle - \langle X^\eta|(I_0^{-1})I_1|\Phi_0^\lambda\rangle + \dots = \langle\Phi^\eta|I_0|\Phi_0^\lambda\rangle - \langle\Phi^\eta|I_1|\Phi_0^\lambda\rangle + \dots \,,\end{aligned} \tag{125}$$

from which one sees that the corrections to the transport coefficient are given by

$$\Xi_1^{\eta,\lambda} = -\langle\Phi_0^\eta|I_1|\Phi_0^\lambda\rangle \,. \tag{126}$$

6.1 Diffusion

In the absence of a temperature gradient, the flux density of ^{3}He atoms in the frame moving with the phonons is given by

$$n_3\mathbf{u} = -\Xi_{cc}\boldsymbol{\nabla}(\mu_3 - \epsilon_0) = -\Xi\frac{T}{n_3}\boldsymbol{\nabla}n_3, \tag{127}$$

and therefore Ξ_{cc} is related to the diffusion coefficient by

$$\Xi_{cc} = Dn_3/T. \tag{128}$$

One then sees from Eq. (125) that the fractional change in the diffusion coefficient due to recoil is

$$\frac{D_1(T)}{D_0(T)} \simeq -\frac{\langle\Phi_0^c|I_1|\Phi_0^c\rangle}{\langle\Phi_0^c|I_0|\Phi_0^c\rangle}. \tag{129}$$

This ratio is precisely that evaluated in the Appendix of Ref. [6], and therefore

$$\frac{D(T)}{D_0(T)} \simeq 1 + \frac{T}{m^*s^2}\left(\frac{100\pi^2 - 198}{33}\frac{\tilde{J}}{J} - 1\right) \,, \tag{130}$$

where $\tilde{J} = 4A^2/3 - 4AB/3 + 8B^2/15 \approx 3.30$ and the coefficient of T/m^*s^2 is 35.5.

The reason that the calculation of recoil corrections is independent of x_3 is that the solution to the ^{3}He Boltzmann equation is $w(p) =$ constant: for the case of low x_3 considered in this paper, this is the exact solution for ^{3}He–phonon scattering when ^{3}He-^{3}He scattering is negligible. This is not altered at higher x_3 because $\Phi(p) \propto p_z$ is still a solution of the Boltzmann equation when ^{3}He-^{3}He collisions are included, since they conserve the total ^{3}He momentum.

An alternative approach to calculating recoil corrections is to start from the standard variational expression for transport coefficients [9], which in the case of diffusion is

$$\frac{Dn_3}{T} \geq \frac{\langle X^c|\Phi\rangle^2}{\langle\Phi|I|\Phi\rangle}, \tag{131}$$

for an arbitrary form of the function Φ. In particular, if one chooses $\Phi = \Phi_0^c$ one has

$$\frac{Dn_3}{T} \geq \frac{\langle X^c|\Phi_0^c\rangle^2}{\langle\Phi_0^c|I|\Phi_0^c\rangle}, \tag{132}$$

which recovers the exact results for the D_0 and D_1 but also has terms of higher order in k^2.

6.2 Thermal diffusion

We turn now to Ξ^{hc}, which describes the heat flow induced by a relative motion of phonons and ^{3}He atoms (and also the relative motion of phonons and ^{3}He induced by a temperature gradient). To first order in T/m^*s^2 this quantity vanishes since $\langle\Phi_0^h|I_0|\Phi_0^c\rangle$ vanishes. This is because $\Phi^c(p) \propto p_z$ is an eigenstate of I_0 and therefore $I_0|\Phi_0^c\rangle \propto |\Phi_0^c\rangle$. As shown in Sec. 4, the solution for the Boltzmann equation for response to a temperature gradient is proportional to $(p^2/2m^* - 5T/2)p_z$ to leading order in T/m^*s^2 and consequently $\langle\Phi_0^h|I_0|\Phi_0^c\rangle \propto \langle\Phi^h|\Phi^c\rangle$ because of the orthogonality relation for Sonine polynomials for $n = 0$ and $n = 1$, $\int_0^\infty dx x^4 \exp(-x^2/2)(x^2 - 5) = 0$. The leading contributions to Ξ^{hc} are therefore of second order in T/m^*s^2, while those to Ξ^{cc} are of first order. In addition, Ξ^{hc} has an extra factor of T compared with Ξ^{cc} because of the extra factor $p^2/2m^* - 5T/2$, and therefore as noted in Sec. (4), the magnitude of χ is $\sim n_3T/m^*s^2$.

In Secs. 2.2 and 2.3 we saw that χ usually enters in combinations such as $n_3 + \chi$, $S_3 + \chi$, or $S_{ph} - \chi$. Therefore, under the conditions of the nEDM proposed experiment, we expect the nonzero value of χ to have little effect and we shall not evaluate the prefactor.

7 Results

We now summarize the main results of this paper in graphical form. Over the full range of ^{3}He concentrations in the non-degenerate regime, from that of the Lamoreaux et al. experiment [4] ($x_3 \sim 10^{-3}$) to that of the proposed nEDM experiment [3] ($x_3 \lesssim 10^{-10}$), the diffusion constant, D, Eq. (84), is independent of x_3, even taking the recoil corrections of Sec. 6 into account. As Fig. 1 shows, its basic T^{-7} temperature dependence is slightly modified by the recoil corrections. As expected, the phonon and ^{3}He thermal conductivities (Eqs. (36) and (89), respectively) depend on both T and x_3. As shown in Fig. 2,



Fig. 1 Diffusion constant multiplied by T^7, as a function of temperature. The results include recoil corrections and were obtained from numerical integration using Eq. (132). We note that the expression (130), which includes only the contributions D_0 and D_1, gives results that lie within 3% of this result over the range shown.

at low ^{3}He concentrations, the phonon thermal conductivity is dominated by phonon viscosity; at the highest concentrations considered here, K_{ph} is simply proportional to $1/x_3$ because of phonon scattering from ^{3}He. On the other hand, the ^{3}He thermal conductivity grows linearly with increasing x_3 at low concentrations before becoming dominated by ^{3}He-^{3}He scattering at concentrations of about $x_3 \sim 10^{-6}$. At all concentrations considered here, the ^{3}He contribution to the transport of heat is $\lesssim 1\%$ that of the phonons. We note that we have not taken into account the effects of geometry on the mean free paths of phonons and ^{3}He, especially important at low temperatures and low concentrations, respectively, in this treatment but will do so in Ref. [12].

8 Summary

In this paper we have calculated the transport properties of ^{3}He in superfluid ^{4}He in the non-degenerate regime $x_3 \lesssim 10^{-3}$ for $T \lesssim 0.6$ K where phonons are the dominant ^{4}He excitations. These calculations are relevant to previous transport measurements at relatively high concentrations [4,7], to the range of natural concentration, $x_3 \sim 10^{-6}$ [8], as well as to the low concentrations expected in the neutron EDM experiment [3]. We began by considering particle number, momentum and energy conservation, including the effects of phonon viscosity. The time evolution of the relative velocity $\mathbf{u} = \mathbf{v}_3 - \mathbf{v}_{ph}$ is given by Eq. (21); the relative velocity is driven by both gradients of the chemical potentials and temperature, and includes the dissipative effects of ^{3}He-phonon



Fig. 2 Phonon and ^{3}He thermal conductivities from Eqs. (36) and (89), respectively, at $T = 0.25$, 0.35 and 0.45 K. Note that the scale of phonon thermal conductivities is 10^5 larger than the scale of ^{3}He thermal conductivities. The phonon thermal conductivity is calculated using the phonon mean free path, Eq. (39), and $k_x^2 = 8/R^2$, with $R = 15$ cm (approximately 10 times the phonon mean free path at $T = 0.25$ K). Both K_{ph} and K_3 are calculated using the recoil correction obtained from numerical integration as described in the caption of Fig. 1

scattering and phonon viscosity. We show explicitly that the entropy generation rate is positive definite. The energy currents lead us to identify the thermal conductivities of the ^{3}He (K_3, Eq. (24)) and of the phonons (K_{ph}, Eq. (36)). In addition we also identify the ordinary and "thermoelectric" diffusion coefficients D and χ, Eq. (25), as well as the Onsager reciprocity relation connecting the ^{3}He heat current and the thermoelectric diffusion coefficient.

Before calculating the transport coefficients in a Boltzmann equation framework, we discuss in detail in Sec. 3 the various scattering rates in the problem. Whereas the ^{3}He-^{3}He scattering rate is much larger than the ^{3}He-phonon rate at the high concentrations of the Lamoreaux, et al. [4] and Rosenbaum, et al. [7] measurements, at $x_3 \sim 10^{-6}$, they are roughly equal, and at lower concentrations, the scattering of ^{3}He by phonons (or walls, depending on the geometry [12]) dominates. In treating phonon-phonon scattering we use the fact that phonons propagating in a given direction in momentum are in local equilibrium, owing to the very large small angle scattering rate. The phonon viscosity arises from scattering of phonons through large angles, either in a single large angle event or in a sequence of small angle scatterings. When effects of phonon viscosity are important, it is necessary to consider the effects of phonons scattering from the walls, the only appearance of geometry in the present paper.

In Secs. 4 and 5 of the paper, we calculate the transport coefficients starting from the Boltzmann equations for the evolution of the ^{3}He and phonon dis-

tribution functions. The ^{3}He-phonon scattering amplitude is well established by measurements from the 1960s [24]-[26]. For the high ^{3}He concentrations of Refs. [4,7], where ^{3}He-^{3}He scattering keeps the ^{3}He in equilibribum, we can ignore the energy transfer relative to the momentum transfer, k, in the ^{3}He-phonon scattering. However, in general, it is necessary to consider both. The leading effects appear at order k^2, for which the ^{3}He Boltzmann equation is effectively a Fokker-Planck equation, but there are important contributions at order k^4, which we describe as recoil corrections (see Sec. 6). We find that the solutions of the Fokker-Planck equation for the ^{3}He distribution function are exactly Sonine polynomials, familiar from calculations of classical transport coefficients of one component gases. As we calculate, the diffusion constant varies approximately as T^{-7}, is independent of x_3, and, including the recoil corrections, has a leading coefficient approximately 70% as large as that found in Ref. [4]. The thermal transport is dominated by that of the phonons for the full range of x_3; for $x_3 \gtrsim 10^{-6}$ ^{3}He-phonon scattering reduces K_{ph} below its low x_3 value where it is determined by phonon viscosity. The ^{3}He thermal conductivity is limited by ^{3}He-^{3}He scattering for $x_3 \gtrsim 10^{-6}$ but is, in any case, much smaller than K_{ph}. Finally, the thermoelectric coefficient, χ, because of its "off-diagonal" nature, vanishes to lowest order in k^2 due to the orthogonality of the Sonine polynomials; the k^4 corrections are furthermore negligible.

References

1. Piet Hein, *Grooks 1* (Doubleday & Co., New York, 1969).
2. See, e.g., G. Baym, C. J. Pethick, *Landau Fermi Liquid Theory: Concepts and A pplications* (J. Wiley and Sons, N. Y.,1991)
3. R. Golub, S. K. Lamoreaux, Phys. Rep. **237**, 1 (1994); S. K. Lamoreaux, R. Golub, J. Phys G.: Nucl. Part. Phys. **36**, 104002 (2009)
4. S. K. Lamoreaux, G. Archibald, P. D. Barnes, W. T. Buttler, D. J. Clark, M. D. Cooper, M. Espy, G. L. Greene, R. Golub, M. E. Hayden, C. Lei, L. J. Marek, J.-C. Peng, S. Penttila, Europhys. Lett. **71**, 724 (2002)
5. R. M. Bowley, Europhys. Lett. **58**, 725-729 (2002)
6. G. Baym, D. H. Beck, C. J. Pethick, Phys. Rev B **88**, 014512 (2013)
7. R. L. Rosenbaum, J. Landau, Y. Eckstein, J. Low Temp. Phys. **16**, 131 (1974)
8. D. Beck et al., nEDM collaboration, to be published.
9. E.M. Lifshitz and L.P. Pitaevskii, *Physics Kinetics* (Pergamon Press, Oxford, 1981), Sec. 10
10. L. D. Landau and E. M. Lifshitz, *Fluid Mechanics* (Pergamon Press, Oxford, 1959), Ch. VI
11. The expansion in the parameter $k/\sqrt{m^*T}$ is essentially one in small angle scattering. The ^{3}He-phonon collision term is thus similar to a Fokker-Planck collision term, to which Sonine polynomials have previous been applied,, e.g., K. Abe, Y. Ushimi, Phys. Fluids **19**, 2047 (1976).
12. G. Baym, D. H. Beck, C. J. Pethick, to be published.
13. I.M. Khalatnikov, V.N. Zharkov, J. Exptl. Theoret. Phys. (USSR) **32**, 1108 (1957) [Engl. transl. Soviet Physics JETP **5**, 95 (1957)]
14. I.M. Khalatnikov, *Introduction to the Theory of Superfluidity*, (W.A. Benjamin, New York, 1965), Chs. 24, 25
15. G. Baym, C. Ebner, Phys. Rev. **164**, 235 (1967)
16. Here, and in experiments determining heat transport in superfluid helium, the $\mathbf{v}_{ph} - \mathbf{v}_s$ achieved depends on the ambient temperature gradient.

17. Ref. [14], pp. 65, 133
18. G. Baym, W. F. Saam, Phys. Rev. **171**, 172 (1968)
19. Compare with Eq. (59.6) of Ref. [10], with the identifications $\mathbf{i} = m_3 n_3 (1-y)\mathbf{u}$ and $\mathbf{q} - \mu\mathbf{i} = \mathbf{Q}_3 - T\sigma\mathbf{u}$.
20. Ref. [10]. Eq. (59.6).
21. D. Benin, H.J. Maris, Phys. Rev. B **18**, 3112 (1978)
22. H. J. Maris, Rev. Mod. Phys. **49**, 341 (1977)
23. D. Greywall, Phys. Rev. B **23**, 2152 (1981), esp. p. 2164 first column near bottom.
24. C. Boghosian, H. Meyer, Phys. Lett. **A25**, 352 (1967)
25. G. E. Watson, J. D. Reppy, R. C. Richardson, Phys. Rev. **188**, 384 (1969)
26. B. M. Abraham, C. G. Brandt, Y. Eckstein, J. Munarin, G. Baym, Phys. Rev. **188**, 309 (1969)
27. In Ref. [6] we wrote the effective ^{3}He-phonon scattering rate as Γ/p^2; we emphasize that when the energy transfer in ^{3}He-phonon collisions is taken into account, the scattering rate is independent of p and has the value $\Gamma/3m^*T$, which is the value of Γ/p^2 for p^2 replaced by its thermal average $3m^*T$.
28. Three phonon Landau damping and the Beliaev process, although they involve phonons alone, can affect the rate at which momentum is transferred from phonons to ^{3}He. These processes conserve the total momentum and energy flux of the phonons. However, the cross section for scattering of phonons by ^{3}He atoms is strongly dependent on the phonon momentum and therefore the total rate at which momentum is transferred from phonons to ^{3}He depends on the details of the phonon distribution, not just the total momentum of the phonons.